

# A NOTE ON ANALYTICAL REPRESENTABILITY OF MAPPINGS INVERSE TO INTEGRAL OPERATORS M.I.OSTROVSKII

**Abstract.** The condition onto pair  $(F, G)$  of function Banach spaces under which there exists a integral operator  $T : F \rightarrow G$  with analytic kernel such that the inverse mapping  $T^{-1} : \text{im} T \rightarrow F$  does not belong to arbitrary a priori given Borel (or Baire) class is found.

We begin with recalling some definitions [4, §31.IX]. Let  $X$  and  $Y$  be metric spaces. By definition, the family of *analytically representable* mappings is the least family of mappings from  $X$  to  $Y$  containing all continuous mappings and being closed with respect to passage to pointwise limit.

This family is representable as an union  $\cup_{\alpha \in \Omega} \Phi_\alpha$ , where  $\Omega$  is the set of all countable ordinals and  $\Phi_\alpha$  are defined in the following way.

1. The class  $\Phi_0$  is the set of all continuous mappings.
2. The class  $\Phi_\alpha$  ( $\alpha > 0$ ) consists of all mappings which are limits of convergent sequences of mappings belonging to  $\cup_{\xi < \alpha} \Phi_\xi$ .

A mapping  $f : X \rightarrow Y$  is called of  $\alpha$  *Borel class* if the set  $f^{-1}(F)$  is a Borel set of multiplicative class  $\alpha$  for every closed subset  $F \subset Y$ .

It is known [1], [11], if  $Y$  is a separable Banach space, then the class  $\Phi_\alpha$  coincides with the  $\alpha$  Borel class for finite  $\alpha$  and with  $\alpha + 1$  Borel class for infinite  $\alpha$ . In addition [12], if  $Y$  is a Banach space, then the class of all regularizable mappings from  $X$  to  $Y$ , important in the theory of improperly posed problems coincides with  $\Phi_1$ .

The present paper is devoted to the following problem. Let  $F$  and  $G$  be Banach function spaces on  $[0, 1]$  and  $T : F \rightarrow G$  be an injective integral operator with analytic kernel. What class of analytically representable functions can the mapping  $T^{-1} : TF \rightarrow F$  belong to?

Let us give some clarifications.

1. By analytic kernel we mean the mapping  $K : [0, 1] \times [0, 1] \rightarrow \mathbf{C}$  with the following property: For some open subsets  $\Gamma$  and  $\Delta$  of  $\mathbf{C}$  such that  $[0, 1] \subset \Gamma$  and  $[0, 1] \subset \Delta$ , there exists an analytic continuation of  $K$  to  $\Gamma \times \Delta$ .
2. If the spaces are real then we consider mapping  $K$  taking real values on  $[0, 1] \times [0, 1]$ .

The main result of the present paper states that for wide class of pairs  $(F, G)$  and every countable ordinal  $\alpha$  there exists an injective integral operator  $T : F \rightarrow G$  with an analytic kernel such that  $T^{-1} : TF \rightarrow F$  does not belong to  $\Phi_\alpha$ .

This result is a generalization of the Menikhes' result [6]. The last states that there exists an integral operator from  $C(0, 1)$  to  $L_2(0, 1)$  with infinite differentiable kernel and nonregularizable inverse. A.N.Plichko [9] generalized this result onto wide classes of function spaces. The relations of our results with the results of [9] will be discussed in remark 3 below.

We use standard Banach space terminology and notation, as may be found in [5].

For a subset  $A$  of a Banach space  $X$  by  $\text{cl}A$ ,  $\text{lin}A$  and  $A^\perp$  we shall denote the closure of  $A$  in the strong topology, the linear span of  $A$  and  $\{x^* \in X^* : (\forall x \in A)(x^*(x) = 0)\}$  respectively. For a subset  $A$  of  $X$  by  $A^\top$  we shall denote  $\{x \in X : (\forall x^* \in A)(x^*(x) =$

0)}. By  $w^* - \lim$  we shall denote the limit in the weak\* topology.

In the present paper we restrict ourselves to the case when  $F$  is a separable Banach function space on the closed interval  $[0,1]$  continuously and injectively embedded into  $L_1(0,1)$ . It is clear that classical separable function spaces satisfy this condition. The case of nonseparable spaces continuously and injectively embedded into  $L_1(0,1)$  is discussed in remark 4 below.

Let  $\Gamma$  be a bounded open subset of the complex plane such that  $\Gamma \supset [0,1]$ . Let  $f : \Gamma \rightarrow \mathbf{C}$  be an analytic function continuous on the closure of  $\Gamma$ . (If we consider real spaces then we assume in addition that  $f$  takes real values on the real axis.) Function  $f$  generates a continuous functional on  $L_1(0,1)$  by means of the formula

$$(f, x) = \int_0^1 f(t)x(t)dt.$$

Since  $F$  is continuously embedded into  $L_1(0,1)$  then  $f$  generates a continuous functional on  $F$ . Let us denote by  $U$  the subset of  $F^*$  consisting of all functionals of such type. Let  $M = \text{cl}U$ . It is clear that  $M$  is a closed subspace of  $F^*$ .

Every element of  $F$  may be considered as a functional on  $M$ . The corresponding embedding of  $F$  into  $M^*$  we denote by  $H$ .

**Remark 1.** The subspace  $M$  is a total subspace of  $F^*$ . It follows from the following facts: 1)  $F$  is injectively embedded into  $L_1(0,1)$ ; 2) the set of functionals generated by polynomials is total in  $(L_1(0,1))^*$ .

The main result of the present paper is the following

**THEOREM 1.** Let  $\text{cl}(H(F))$  be of infinite codimension in  $M^*$  and let  $G$  contains a linearly independent infinite sequence of functions which have analytic continuations onto some open region  $\Delta$  of the complex plane such that  $\Delta \supset [0,1]$ . Then for every countable ordinal  $\alpha$  there exists a linear continuous injective integral operator  $T : F \rightarrow G$  with analytic kernel such that  $T^{-1} : TF \rightarrow F$  does not belong to  $\Phi_\alpha$ .

Let us introduce necessary definitions. Let  $X$  be a Banach space. *Weak\* sequential closure* of a subset  $V$  of  $X^*$  is defined to be the set of all limits of weak\* convergent sequences from  $V$ . Weak\* sequential closure will be denoted by  $V_{(1)}$ . For ordinal  $\alpha$  *weak\* sequential closure of order  $\alpha$*  of a subset  $V$  of  $X^*$  is defined by the equality  $V_{(\alpha)} = \cup_{\beta < \alpha} (V_{(\beta)})_{(1)}$ .

**THEOREM 2** (A.N.Plichko [10]). Let  $X$  be a separable Banach space and  $Y$  be an arbitrary Banach space. Let  $T : X \rightarrow Y$  be a continuous linear injective operator. Let  $T^{-1} : TX \rightarrow X$  belongs to  $\Phi_\alpha$ . If  $\alpha$  is finite then  $(T^*Y^*)_{(\alpha)} = X^*$ . If  $\alpha$  is infinite then  $(T^*Y^*)_{(\alpha+2)} = X^*$ . Conversely, if  $\alpha$  is finite and  $(T^*Y^*)_{(\alpha)} = X^*$  or if  $\alpha$  is infinite and  $(T^*Y^*)_{(\alpha+1)} = X^*$ , then the mapping  $T^{-1} : TX \rightarrow X$  belongs to  $\Phi_\alpha$ .

The proof of Theorem 1 is based on Theorem 2 and the following proposition.

**Proposition.** Let  $F$  and  $M$  satisfy the assumptions of Theorem 1. Then for every countable ordinal  $\beta$  there exists a subspace  $K$  of  $M$  such that  $K_{(\beta)} \neq F^*$ .

**Proof.** Using the result due to Davis and Johnson [2, p. 360] we find a weak\* null sequence  $\{u_n\}_{n=1}^\infty$  in  $M$  and a bounded sequence  $\{v_k\}_{k=1}^\infty$  in  $F^{**}$  such that for some partition  $\{I_k\}_{k=1}^\infty$  of the set of natural numbers onto the pairwise disjoint infinite subsets

we shall have

$$v_k(u_n) = \begin{cases} 1, & \text{if } n \in I_k; \\ 0, & \text{if } n \notin I_k. \end{cases}$$

Using the arguments from the proof of Theorem III.1 in [3] we select a weak\* basic subsequence  $\{u_{n(i)}\}_{i=1}^\infty \subset \{u_n\}_{n=1}^\infty$  such that the intersection  $I_k \cap \{n(i)\}_{i=1}^\infty$  is infinite for every  $k \in \mathbf{N}$ . Passing if necessary to a subsequence and renumbering  $\{u_{n(i)}\}_{i=1}^\infty$  we obtain the sequence  $\{y_j\}_{j=0}^\infty$  such that

$$v_k(y_j) = \begin{cases} 1, & \text{if } j \text{ can be represented in the form } j = n(n+1)/2 + k \text{ with } k \leq n; \\ 0, & \text{if it is not the case.} \end{cases}$$

Definition of the weak\* basic sequence implies that the space  $Z = F/(\{y_j\}_{j=0}^\infty)^\top$  has the basis  $\{z_j\}_{j=0}^\infty$  such that  $y_j(z_k) = \delta_{j,k}$ . It is easy to see that the sequence  $\{z_j\}_{j=0}^\infty$  is bounded away from zero and that the set  $\{\sum_{i=j}^k z_{i(i+1)/2+j}\}_{j=0, k=j}^{\infty, \infty}$  is bounded.

We may assume without loss of generality that  $\|z_i\| \leq 1$  for every  $i \in \mathbf{N}$ . Using the arguments from the proof of lemma 1 in [7] we find a subspace  $N$  of  $\text{cl}(\text{lin}\{y_j\}_{j=0}^\infty)$  and a bounded sequence  $\{h_n\}_{n=1}^\infty$  in  $Z^{**}$  such that the following conditions are satisfied:

a) If a weak\* convergent sequence  $\{x_m^*\}_{m=1}^\infty$  is contained in  $N_{(\gamma)}$  for some  $\gamma < \beta$  and  $x^* = w^* - \lim_{m \rightarrow \infty} x_m^*$  then

$$h_n(x^*) = \lim_{m \rightarrow \infty} h_n(x_m^*)$$

for every  $n \in \mathbf{N}$ .

b) There exists a collection  $\{x_{n,m}^*\}_{n=1, m=1}^\infty$  of vectors of  $N_{(\beta)}$  such that for every  $k, n \in \mathbf{N}$  we have

$$\begin{aligned} w^* - \lim_{m \rightarrow \infty} x_{n,m}^* &= 0; \\ (\forall m \in \mathbf{N})(h_k^*(x_{n,m}^*) &= \delta_{k,n}). \end{aligned}$$

Let  $\{s_k^*\}_{k=1}^\infty$  be a total (over  $F$ ) sequence in  $M$ . Let  $c_1 = \sup_n \|h_n\|$ . Let  $\nu_n > 0$  ( $n \in \mathbf{N}$ ) be such that  $\sum_{n=1}^\infty \nu_n < 1/(2c_1)$ .

We shall identify  $Z^*$  with its natural image in  $F^*$ . Without loss of generality we may assume that  $Z^* \neq F^*$ .

Let us denote by  $g_n$  ( $n \in \mathbf{N}$ ) norm-preserving extensions of  $h_n$  onto  $F^*$ . Let us introduce the operator  $R : F^* \rightarrow F^*$  by the equality

$$R(x^*) = x^* + \sum_{n=1}^\infty \nu_n g_n(x^*) s_n^*.$$

It is easy to check that  $\|R - I\| \leq 1/2$  (where  $I$  is the identity operator), hence  $R$  is an isomorphism.

Let  $K = R(N)$ . In the same way as in [7] we prove that for every  $\gamma \leq \beta$  we have

$$K_{(\gamma)} = R(N_{(\gamma)}). \tag{1}$$

Since  $R$  is an isomorphism then relations  $K_{(\beta)} = R(N_{(\beta)})$  and  $N_{(\beta)} \subset Z^* \neq F^*$  imply that  $K_{(\beta)} \neq F^*$ . At the same time (1) and b) imply that  $R(x_{n,m}^*) = x_{n,m}^* + \nu_n s_n^* \in K_{(\beta)}$  and therefore  $s_n^* \in K_{(\beta+1)}$ . By the totality of  $\{s_n^*\}$  it follows that the subspace  $K \subset F^*$  is total. The proof of the proposition is complete.

Proof of Theorem 1. Using the proposition for  $\beta = \alpha + 2$  we find a subspace  $K$  of  $M$  such that  $K_{(\alpha+2)} \neq F^*$ . Using well-known arguments (see e.g. [5, p. 43,44]) we find a fundamental minimal sequence  $\{f_i\}_{i=1}^\infty$  in the space  $F$  such that its biorthogonal functionals  $\{f_i^*\}_{i=1}^\infty$  are total and are contained in  $K$ . It is easy to see (see e.g. [8]) that there exists an isomorphism  $S : F \rightarrow F$  such that functionals  $S^* f_i^* (i \in \mathbf{N})$  are contained in  $U$ . Let us introduce the notation  $g_i^* = S^* f_i^* (i \in \mathbf{N})$ . It is clear that

$$(\text{cl}(\text{lin}(\{g_i^*\}_{i=1}^\infty)))_{(\alpha+2)} \neq F^*. \quad (2)$$

Reducing if necessary the domain  $\Delta$  we may assume that  $\Delta$  is bounded and that  $G$  contains a linearly independent sequence of functions which have analytic continuations onto  $\Delta$  which are continuous on the closure of  $\Delta$ . Using standard biorthogonalization procedure (see e.g. [5, p. 43,44]) we find a minimal sequence  $\{q_i\}_{i=1}^\infty$  in  $G$  such that there exist analytic continuations  $\{\tilde{q}_i\}_{i=1}^\infty$  of  $\{q_i\}_{i=1}^\infty$  to  $\Delta$  which are continuous on the closure of  $\Delta$ .

Let us introduce numbers

$$a_i = \max\{||q_i||_G, \sup_{t \in \Delta} |\tilde{q}_i(t)|\}, (i \in \mathbf{N}).$$

Let  $r_i (i \in \mathbf{N})$  be functions analytic in  $\Gamma$  and continuous on the closure of  $\Gamma$  such that functionals generated by them on  $F$  coincide with  $g_i^* (i \in \mathbf{N})$ . Let us introduce the numbers

$$b_i = \max\{||g_i^*||_{F^*}, \sup_{t \in \Gamma} |r_i(t)|\}, (i \in \mathbf{N}).$$

Let us introduce the operator  $T : F \rightarrow G$  by the equality

$$T(f) = \sum_{i=1}^\infty g_i^*(f) q_i / (2^i a_i b_i). \quad (3)$$

This operator may be considered as an integral operator with the kernel

$$K(t_1, t_2) = \sum_{i=1}^\infty r_i(t_1) \tilde{q}_i(t_2) / (2^i a_i b_i).$$

By the definitions of  $a_i$  and  $b_i$  it follows that this series converges to a function analytic in  $\Gamma \times \Delta$ .

Operator  $T$  is injective since the sequence  $\{g_i^*\}_{i=1}^\infty$  is total and the sequence  $\{q_i\}_{i=1}^\infty$  is minimal. It is easy to see that  $T^* G^* \subset \text{cl}(\text{lin}(\{g_i^*\}_{i=1}^\infty))$ . By (2) we obtain  $(T^* G^*)_{(\alpha+2)} \neq F^*$ . By Theorem 2 it follows that  $T^{-1}$  does not belong to  $\Phi_\alpha$ . The theorem is proved.

**Remark 2.** It is easy to see that if  $F$  and  $G$  are infinite dimensional function spaces and  $T : F \rightarrow G$  is an injective integral operator with an analytic kernel then  $G$  satisfies the condition of Theorem 1.

**Remark 3.** If the subspace  $M \subset F^*$  is norming then the infinite codimension of  $\text{cl}(H(F))$  in  $M^*$  is equivalent to the infinite codimension of  $\text{lin}(M^\perp \cup F)$  in  $F^{**}$ . Therefore Theorems 1 and 2 of [9] may be obtained using the arguments of the present paper. Furthermore, it follows that many of the spaces considered in [9] satisfy the

condition of Theorem 1 of the present paper. In particular it is the case for classical nonreflexive spaces.

**Remark 4.** Let  $F$  be a nonseparable function space, which is continuously injectively embedded into  $L_1(0, 1)$ , and  $G$  is the space satisfying the condition of Theorem 1. Let  $\{g_i^*\}_{i=1}^\infty$  be a total sequence in  $F^*$  such that functionals  $g_i^*(i \in \mathbf{N})$  are represented by functions analytic in the same open subset containing  $[0, 1]$ . Let  $\{q_i\}_{i=1}^\infty$  be the sequence introduced in the proof of Theorem 1. Let us introduce the operator  $T : F \rightarrow G$  by (3). The operator  $T$  is an injective integral operator with an analytic kernel. It is clear that the image of  $T$  is separable. Therefore the mapping  $T^{-1} : TF \rightarrow F$  is not analytically representable since the image of the separable space under the action of analytically representable mapping is separable.

#### REFERENCES

1. Banach S. Über analytisch darstellbare Operationen in abstrakten Räumen, Fund. Math. 17 (1931), 283–295.
2. Davis W.J., Johnson W.B. Basic sequences and norming subspaces in non-quasi-reflexive Banach spaces, Israel J. Math. 14 (1973), 353–367.
3. Johnson W.B., Rosenthal H.P. On  $w^*$ -basic sequences and their applications to the study of Banach spaces, Studia Math. 43 (1972), 77–92.
4. Kuratowski K. Topology, v. I, New York, Academic Press, 1966.
5. Lindenstrauss J., Tzafriri L. Classical Banach spaces, v. I, Berlin, Springer-Verlag, 1977.
6. Menikhes L.D. On the regularizability of mappings inverse to integral operators, Doklady AN SSSR, 241 (1978), no. 2, 282–285 (Russian).
7. Ostrovskii M.I. Total subspaces with long chains of nowhere norming weak\* sequential closures, Note Mat., to appear.
8. Ostrovskii M.I. Regularizability of inverse linear operators in Banach spaces with bases, Sibirsk. Mat. Zh. 33 (1992), no. 3, 123–130 (Russian).
9. Plichko A.N. Non-norming subspaces and integral operators with non-regularizable inverses, Sibirsk. Mat. Zh. 29 (1988), no. 4, 208–211 (Russian).
10. Plichko A.N. Weak\* sequential closures and  $B$ -measurability, private communication.
11. Rolewicz S. On the inversion of non-linear transformations, Studia Math. 17 (1958), 79–83.
12. Vinokurov V.A. Regularizability and analytical representability, Doklady AN SSSR 220 (1975), no. 2, 269–272 (Russian).